

PHENIX Studies of the Scaling Properties of Elliptic Flow at RHIC

A. Taranenko for the PHENIX Collaboration[‡]

Department of Chemistry, SUNY , Stony Brook NY 11794

E-mail: arkadij@rcf.rhic.bnl.gov

Abstract. Recent PHENIX elliptic flow (v_2) measurements for identified particles produced in Au+Au and Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV are presented and compared to other RHIC measurements. They indicate universal scaling of v_2 compatible with partonic collectivity leading to the flow of light, strange and heavy quarks with a common expansion velocity field.

1. Introduction

Universal scaling of elliptic flow (v_2) has been recently observed at RHIC [1, 2, 3]. That is, for a broad range of particle species, v_2/n_q vs. KE_T/n_q scales to a single function; here, n_q and KE_T are the number of valence quarks ($n_q = 2, 3$ for mesons and baryons respectively) and the transverse kinetic energy of the particle. This observation has been interpreted as evidence that transverse expansion of the matter produced in energetic RHIC collisions, occurs during a phase dominated by partonic collectivity.

In this contribution we further demonstrate; (i) the partonic origin of v_2 via scaling tests for the ϕ meson; (ii) essentially full thermalization for the charm quark via scaling tests for D meson v_2 ; and (iii) validate universal scaling of v_2 in Au+Au and Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV.

2. Further tests for universal scaling of elliptic flow at RHIC

Differential v_2 measurements for multi-strange hadrons (ϕ, Ξ, Ω) allow us to address the important question of how the existence of a hadronic phase affects v_2 . These particles are to have a small hadronic cross section for interaction with non-strange hadrons. Consequently, if elliptic flow was established in a phase involving hadrons interacting with their standard hadronic cross sections (post-hadronization), one would expect v_2 for the multi-strange hadrons to be significantly smaller than that for other hadrons (e.g. p, K, π). The left panel of Fig. 1 compares PHENIX v_2 results for the ϕ meson, $\pi^\pm, K^\pm$

[‡] For the full list of PHENIX authors and acknowledgements, see Appendix 'Collaborations' of this volume.

and $(\bar{p})p$ from 20-60% central Au+Au collisions; they show that, despite its mass which is comparable to that for the p and Λ , its $v_2(K E_T)$ values are similar to those for the lighter mesons (π and K) whose hadronic re-scattering cross sections are expected to be large. The right panel of Fig. 1 shows the positive results from a validation test for universal scaling of the ϕ meson v_2 . The recent high-statistics preliminary STAR results for Ω and Ξ produced in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [4] show a similar universal scaling [3]. These results suggest that the v_2 values for hadrons comprised of u, d and s quarks develop in the partonic phase and the existence of a hadronic phase does not lead to their modification.




Figure 1: v_2 vs. KE_T . and v_2/n_q vs. KE_T/n_q for π, K, p and ϕ mesons detected in semi-central (20-60%) Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV.

Further insight into properties of the produced medium at RHIC can be obtained from the study of elliptic flow of particles comprised of heavy quarks (charm or bottom). If charm quarks flow like light quarks, it would indicate an unexpectedly strong interaction of charm with the medium. One method of accessing charm v_2 is to measure the v_2 for “non-photonic” electrons [5]. A recent high statistics PHENIX measurements show rather larger elliptic flow for “non-photonic” electrons from Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [5]. The “non-photonic” electrons arise mainly from D meson decay for $p_T < 2\text{GeV}/c$. Therefore, the large heavy flavor electron v_2 indicates that the D meson and consequently the charm quark, do flow with a large v_2 . This suggests that the in-medium interactions are strong and perhaps frequent enough to thermalize the charm quark. If this is indeed the case, then $v_2(KE_T)$ for the D meson should show the same universal scaling presented earlier for other hadrons.

The v_2 values for D mesons can be obtained from the values for non-photonic electrons via detailed simulations, see [5] for details. For these simulations, it is necessary to assume the shape for $v_2(p_T)$ for the D mesons. The left and right panels of Fig. 2 compare the unscaled and scaled results (respectively) for v_2 vs. KE_T for π, K, p and D mesons measured in minimum bias Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. Here, the assumed shape for the p_T dependence of the D meson v_2 is that for the proton (see




Figure 2: v_2 vs. KE_T . and v_2/n_q vs. KE_T/n_q . for $\pi^\pm, K^\pm, (\bar{p})p$ from minimum bias Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [1]. Results for D-meson is shown for the proton-like assumption for the shape of D $v_2(p_T)$ [3, 5]

[3, 5]). This assumption is compatible with a large array of differential v_2 measurements for different “heavy” particle species [1, 3, 6]. Both panels of Fig. 2 show robust scaling; they confirm that the D mesons (for $p_T < 2.0$ GeV/c) and hence, their associated charm quarks, flow with a fluid velocity field similar to that for the other quarks. This suggests that the medium responds as a thermalized fluid and the transport mean free path is small.

If v_2 develops in a phase dominated by partonic collectivity and u, d, s and c quarks flow with a common velocity field, one expects universal scaling for all particle species measured at RHIC. Figure 3(a) shows differential flow measurements $v_2(p_T)$, for several particle species produced at mid-rapidity in central and semi-central Au+Au collisions at $\sqrt{s_{NN}}=200$ GeV; they span essentially the full range of measurements (several hundred data points) at RHIC. The v_2 values for pions ($\pi^\pm, \pi^0$), kaons $K^\pm$, (anti-)protons ($\bar{p}, p$), (anti-)deuterons ($\bar{d}, d$) and the ϕ meson comprise results from the PHENIX [1, 2, 3, 7]. The values for neutral kaons (K^0), lambdas ($\Lambda + \bar{\Lambda}$), cascades ($\Xi + \bar{\Xi}$) and omegas ($\Omega + \bar{\Omega}$) represent results from the STAR [4, 6]. Fig. 3 (b) shows the scaled results ($v_2/n_q \epsilon$ vs KE_T/n_q) obtained from the same data; here ϵ is the integral v_2 of charged hadrons for each of the indicated centrality selections, multiplied by a constant factor $k \sim 3.2$ (i.e. $\epsilon = k \times v_2$) [1, 2, 3]. Recent measurements [1] indicate that $v_2(p_T)/\epsilon$ is independent of centrality and the size of the colliding system as would be expected from a hydrodynamic system. Fig. 3(b) indicates that the relatively complicated dependence of v_2 on centrality, transverse momentum, particle type, etc., for particles produced at mid-rapidity can be scaled to a single function. Fig. 1(c) demonstrates that the same scaling also holds for $\pi^\pm, (K^\pm)$ and $(\bar{p}, p)$ produced at mid-rapidity in Cu+Cu collisions at $\sqrt{s_{NN}}=200$ GeV.




Figure 3: (a) v_2 vs. p_T and (b) $v_2/(n_q \epsilon)$ vs. KE_T/n_q for several centralities and particle species as indicated. (c) $v_2/(n_q \epsilon)$ vs. KE_T/n_q for $\pi^\pm$, $K^\pm$, and $(\bar{p}, p)$ from Cu+Cu and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV

3. Summary

Validation tests for the universal scaling of v_2 at RHIC, suggest that the transverse expansion dynamics leading to elliptic flow are not controlled by ordinary hadrons interacting with their standard hadronic cross sections. Instead, they suggest a pre-hadronization state exhibiting partonic collectivity leading to the flow of light, strange and heavy quarks with a common expansion velocity field.

References

- [1] PHENIX Collaboration, A. Adare *et.al.* nucl-ex/0608033
- [2] PHENIX Collaboration, M. Issah *et.al.* nucl-ex/0604011
- [3] R. A. Lacey, A. Taranenko, nucl-ex/0610029
- [4] STAR Collaboration, M. Oldenburg *et.al.*, *J. Phys. G* **32**, S563 (2006)
- [5] PHENIX Collaboration, S. Sakai *et.al.*, *J. Phys. G* **32**, S551 (2006); PHENIX Collaboration, A. Adare *et.al.* nucl-ex/0611018;
- [6] STAR Collaboration, J. Adams *et.al.* *Phys. Rev. Lett.* **92**, 052302 (2004); J. Adams *et.al.* *Phys. Rev. C* **72**, 014904 (2005), J. Adams *et.al.* *Phys. Rev. Lett.* **95**, 122301 (2005)
- [7] PHENIX Collaboration, S. S. Adler *et.al.* *Phys. Rev. Lett.* **96**, 032302 (2006)